

REFERENCES

1. Oesser, S., et al., Oral administration of (14)C labeled gelatin hydrolysate leads to an accumulation of radioactivity in cartilage of mice (C57/BL). *J Nutr*, 1999. 129(10): p. 1891-5.
2. Oesser, S. and J. Seifert, Stimulation of type II collagen biosynthesis and secretion in bovine chondrocytes cultured with degraded collagen. *Cell Tissue Res*, 2003. 311(3): p. 393-9.
3. Dar, Q.A., et al., Oral hydrolyzed type 1 collagen induces chondroregeneration and inhibits synovial inflammation in murine posttraumatic osteoarthritis. *Osteoarthritis and Cartilage*, 2016. 24: p. S532-S533.
4. Paddon-Jones, D., et al., Amino acid ingestion improves muscle protein synthesis in the young and elderly. *Am J Physiol Endocrinol Metab*, 2004. 286(3): p. E321-8.
5. Moore, D.R., et al., Ingested protein dose response of muscle and albumin protein synthesis after resistance exercise in young men. *Am J Clin Nutr*, 2009. 89(1): p. 161-8.
6. Reidy, P.T., et al., Protein blend ingestion following resistance exercise promotes human muscle protein synthesis. *J Nutr*, 2013. 143(4): p. 410-6.
7. Paddon-Jones, D., et al., Exogenous amino acids stimulate human muscle anabolism without interfering with the response to mixed meal ingestion. *Am J Physiol Endocrinol Metab*, 2005. 288(4): p. E761-7.
8. Clifton, P.M., D. Condo, and J.B. Keogh, Long term weight maintenance after advice to consume low carbohydrate, higher protein diets--a systematic review and meta analysis. *Nutr Metab Cardiovasc Dis*, 2014. 24(3): p. 224-35.
9. Santesso, N., et al., Effects of higher- versus lower-protein diets on health outcomes: a systematic review and meta-analysis. *Eur J Clin Nutr*, 2012. 66(7): p. 780-8.
10. Lejeune, M.P., E.M. Kovacs, and M.S. Westerterp-Plantenga, Additional protein intake limits weight regain after weight loss in humans. *Br J Nutr*, 2005. 93(2): p. 281-9.
11. Weigle, D.S., et al., A high-protein diet induces sustained reductions in appetite, ad libitum caloric intake, and body weight despite compensatory changes in diurnal plasma leptin and ghrelin concentrations. *Am J Clin Nutr*, 2005. 82(1): p. 41-8.
12. Mojtahedi, M.C., et al., The effects of a higher protein intake during energy restriction on changes in body composition and physical function in older women. *J Gerontol A Biol Sci Med Sci*, 2011. 66(11): p. 1218-25.
13. Leidy, H.J., et al., The role of protein in weight loss and maintenance. *Am J Clin Nutr*, 2015.
14. Zdzieblik, D., et al., Collagen peptide supplementation in combination with resistance training improves body composition and increases muscle strength in elderly sarcopenic men: a randomised controlled trial. *Br J Nutr*, 2015. 114(8): p. 1237-45.
15. Hays, N.P., et al., Effects of whey and fortified collagen hydrolysate protein supplements on nitrogen balance and body composition in older women. *J Am Diet Assoc*, 2009. 109(6): p. 1082-7.
16. Paddon-Jones, D., et al., Protein, weight management, and satiety. *Am J Clin Nutr*, 2008. 87(5): p. 1558S-1561S.
17. Stubbs, J., S. Ferres, and G. Horgan, Energy density of foods: effects on energy intake. *Crit Rev Food Sci Nutr*, 2000. 40(6): p. 481-515.
18. Veldhorst, M.A., et al., A breakfast with alpha-lactalbumin, gelatin, or gelatin + TRP lowers energy intake at lunch compared with a breakfast with casein, soy, whey, or whey-GMP. *Clin*

Nutr, 2009. 28(2): p. 147-55.

19. Hochstenbach-Waelen, A., et al., Single-protein casein and gelatin diets affect energy expenditure similarly but substrate balance and appetite differently in adults. *J Nutr*, 2009. 139(12): p. 2285-92.

20. Pesta, D.H. and V.T. Samuel, A high-protein diet for reducing body fat: mechanisms and possible caveats. *Nutr Metab (Lond)*, 2014. 11(1): p. 53.

21. Westerterp-Plantenga, M.S., et al., Satiety related to 24 h diet-induced thermogenesis during

high protein/carbohydrate vs high fat diets measured in a respiration chamber. *Eur J Clin Nutr*, 1999. 53(6): p. 495-502.

22. Whitehead, J.M., G. McNeill, and J.S. Smith, The effect of protein intake on 24-h energy expenditure during energy restriction. *Int J Obes Relat Metab Disord*, 1996. 20(8): p. 727-32.

23. Mikkelsen, P.B., S. Toubro, and A. Astrup, Effect of fat-reduced diets on 24-h energy expenditure: comparisons between animal protein, vegetable protein, and carbohydrate. *Am J Clin Nutr*, 2000. 72(5): p. 1135-41.

24. Darling, A.L., et al., Dietary protein and bone health: a systematic review and meta-analysis. *Am*

J Clin Nutr, 2009. 90(6): p. 1674-92.

25. MacDonell, R., M.W. Hamrick, and C.M. Isales, Protein/amino-acid modulation of bone cell function. *Bonekey Rep*, 2016. 5: p. 827.

26. Daneault, A., Hydrolyzed collagen contributes to osteoblast differentiation in vitro and subsequent bone health in vivo. *Osteoarthritis and Cartilage*, 2014. 22: p. S131.

27. Guillerminet, F., et al., Hydrolyzed collagen improves bone metabolism and biomechanical parameters in ovariectomized mice: an in vitro and in vivo study. *Bone*, 2010. 46(3): p. 827-34.

28. Guillerminet, F., et al., Hydrolyzed collagen improves bone status and prevents bone loss in ovariectomized C3H/HeN mice. *Osteoporos Int*, 2012. 23(7): p. 1909-19.

29. Martin-Bautista, E., et al., A nutritional intervention study with hydrolyzed collagen in prepubertal

spanish children: influence on bone modeling biomarkers. *J Pediatr Endocrinol Metab*, 2011. 24(3-4): p. 147-53.

30. Allegrini, A. and M. Costantini, Gelatine tannate for the treatment of acute diarrhoea in adults.

Journal of Gastrointestinal and Digestive System, 2012. 2.

31. Lazzi, C., et al., Growth promotion of Bifidobacterium species by poultry bone and meat trimming hydrolyzate. *J Food Sci*, 2011. 76(6): p. M392-7.

32. Lazzi, C., et al., Growth promotion of Bifidobacterium and Lactobacillus species by proteinaceous

hydrolysates derived from poultry processing leftovers. *International Journal of Food Science and Technology*, 2013. 48: p. 341-349.

33. Meli, F., et al., Effect of protein hydrolyzates on growth kinetics and aminopeptidase activities of

some Bifidobacterium species. *Anaerobe*, 2013. 22: p. 130-3.

34. Jiang, J.-X., et al., Collagen peptides improve knee osteoarthritis in elderly women: A 6-month

randomized, double-blind, placebo-controlled study. *Agro FOOD Industry Hi Tech*, 2014. 25: p. 19-23.

35. Moskowitz, R.W., Role of collagen hydrolysate in bone and joint disease. *Semin Arthritis Rheum*,

2000. 30(2): p. 87-99.

36. Kumar, S., et al., A double-blind, placebo-controlled, randomised, clinical study on the

- effectiveness of collagen peptide on osteoarthritis. *J Sci Food Agric*, 2015. 95(4): p. 702-7.
37. Clark, K.L., et al., 24-Week study on the use of collagen hydrolysate as a dietary supplement in athletes with activity-related joint pain. *Curr Med Res Opin*, 2008. 24(5): p. 1485-96.
38. Trc, T. and J. Bohmova, Efficacy and tolerance of enzymatic hydrolysed collagen (EHC) vs. glucosamine sulphate (GS) in the treatment of knee osteoarthritis (KOA). *Int Orthop*, 2011. 35(3): p. 341-8.
39. Benito-Ruiz, P., et al., A randomized controlled trial on the efficacy and safety of a food ingredient, collagen hydrolysate, for improving joint comfort. *Int J Food Sci Nutr*, 2009. 60 Suppl 2: p. 99-113.
40. McAlindon, T.E., et al., Change in knee osteoarthritis cartilage detected by delayed gadolinium enhanced magnetic resonance imaging following treatment with collagen hydrolysate: a pilot randomized controlled trial. *Osteoarthritis Cartilage*, 2011. 19(4): p. 399-405.
41. Ribas Fernández, J. and O. Molinero Pérez, Effects of gelatine hydrolysate in the prevention of athletic injuries. *Archivos de Medicina del Deporte*, 1998. 15(66): p. 277-282.
42. Baar, K., Training and Nutrition to Prevent Soft Tissue Injuries and Accelerate Return to Play. *Sports Science Exchange* 2015. 28(142): p. 1-6.
43. Shaw, G., et al., Vitamin C-enriched gelatin supplementation before intermittent activity augments collagen synthesis. *Am J Clin Nutr*, 2017. 105(1): p. 136-143.
44. Minaguchi, J., et al., Effects of ingestion of collagen peptide on collagen fibrils and glycosaminoglycans in Achilles tendon. *J Nutr Sci Vitaminol (Tokyo)*, 2005. 51(3): p. 169-74.
45. Wienecke, E., Performance Explosion in Sports: An Anti-doping Concept: Revolutionary New Findings in the Area of Micronutrient Therapy: Training Continuity, Training Optimization, Injury Prevention Through Personalized Micronutrients. 2011: Meyer & Meyer Verlag.
46. Proksch, E., et al., Oral intake of specific bioactive collagen peptides reduces skin wrinkles and increases dermal matrix synthesis. *Skin Pharmacol Physiol*, 2014. 27(3): p. 113-9.
47. Proksch, E., et al., Oral supplementation of specific collagen peptides has beneficial effects on human skin physiology: a double-blind, placebo-controlled study. *Skin Pharmacol Physiol*, 2014. 27(1): p. 47-55.
48. Asserin, J., et al., The effect of oral collagen peptide supplementation on skin moisture and the dermal collagen network: evidence from an ex vivo model and randomized, placebo-controlled clinical trials. *J Cosmet Dermatol*, 2015. 14(4): p. 291-301.
49. PM, C., et al., An Oral Supplementation Based on Hydrolyzed Collagen and Vitamins Improves Skin Elasticity and Dermis Echogenicity: A Clinical Placebo-Controlled Study. *Clinical Pharmacology & Biopharmaceutics*, 2015. 4(3).
50. Choi, S.Y., et al., Effects of collagen tripeptide supplement on skin properties: a prospective, randomized, controlled study. *J Cosmet Laser Ther*, 2014. 16(3): p. 132-7.
51. Ohara, H., et al., Improvement in the Moisture Content of the Stratum Corneum Following 4 Weeks of Collagen Hydrolysate Ingestion. *Nippon Shokuhin Kagaku Kogaku Kaishi*, 2009. 56(3): p. 137-145.
52. Borumand, M. and S. Sibilla, Daily consumption of the collagen supplement Pure Gold Collagen(R) reduces visible signs of aging. *Clin Interv Aging*, 2014. 9: p. 1747-58.

53. Liang, J., et al., The protective effects of long-term oral administration of marine collagen hydrolysate from chum salmon on collagen matrix homeostasis in the chronological aged skin of Sprague-Dawley male rats. *J Food Sci*, 2010. 75(8): p. H230-8.